



315-253-8408  
Auburn Office

315-781-8502  
Geneva Office

**AS A COURTESY TO OUR PATIENTS KINDLY GIVE US A  
48 HOUR NOTICE TO CANCEL THANK YOU.**

### **PRE-OPERATIVE INSTRUCTIONS**

#### **Before oral surgery and general anesthesia or IV sedation**

Avoid the use of aspirin or aspirin products such as Anacin, Bufferin, or Alka-Seltzer, for at least three days prior to surgery; Tylenol is OK to use instead.

If you have general anesthesia or intravenous sedation, **A RESPONSIBLE PERSON MUST ACCOMPANY YOU, REMAIN IN THE RECEPTION AREA, AND TAKE YOU HOME.** You **must NOT drive** a car or operate machinery for 24 hours after intravenous sedation or general anesthesia. It is also strongly recommended that a responsible person be with the patient at home the first day to assist them as they complete recovery from any anesthetics or sedation.

#### **Please read these additional important pre-operative instructions:**

1. If you chose to have general anesthesia or intravenous sedation, **do not have anything to eat or drink, including water, after Midnight the night before.**
2. If you take medications regularly, you should take them with small sips of water at the regular time. Also, any additional medications, which might be prescribed to you by our office, should be taken with small sips of water as directed.
3. Wear comfortable, loose fitting clothing with short sleeves or sleeves that can easily be rolled up above both elbows.
4. Do not wear jewelry, including any nose, mouth, tongue or other facial piercings or makeup, including nail polish or eye cosmetics. Please tie long hair back.
5. If you wear contact lenses, please remove them prior to your appointment.
6. No smoking the morning of surgery and preferably for at least one week afterwards.
7. Your mouth and teeth should be well cleansed to help avoid infection.
8. Do not ignore a head or chest cold when oral surgery is to be performed. Please *call the office if you have any symptoms, as an appointment change may be necessary.*
9. If you wear dentures, please do not use denture adhesive.